

IFT-01/22

July, 2001

Physics of Extra Dimensions as IR Limit of Four-Dimensional Theories*

Stefan Pokorski¹

¹ *Institute for Theoretical Physics*

Hoza 69, 00-681, Warsaw, Poland

February 1, 2008

Abstract

I discuss the recent work done in collaboration with Chris Hill, Jing Wang and Hsin-Chia Cheng. We construct four-dimensional renormalizable gauge theories which, in their infrared limit, generate the dynamics of gauge interactions in flat extra dimensions. In this construction, the Kaluza-Klein states of extra dimensions arise from having many gauge symmetries in four dimensions. The IR limit is determined by the dynamics of spontaneous breakdown of those symmetries.

*Talk given at the IPPP Workshop on the Phenomenology of Beyond the Standard Model, Durham, Great Britain, 7-10 May 2001, and at Planck2001 “From the Planck Scale to the Electroweak Scale”, La Londe-les-Maures, France, 11-16 May 2001

Non-abelian gauge theories in $4+n$ dimensions have dimensionful coupling constant $g_0 = [M]^{-n/2}$ and are non-renormalizable. They require a cut-off at some scale M_s . Such theories should be viewed as the effective low energy limit of some more fundamental theories, with better UV behaviour.

It is an interesting question how uniquely the physics of a gauge theory in $4+n$ dimensions defines its underlying UV “completion”. One may expect that there exist (universality) classes of theories that have identical behaviour in the infrared but are different above the scale M_s . For instance, it is possible that a given non-abelian gauge theory in $4+n$ dimensions has some stringy UV completion. In ref. [1] we demonstrate a more surprising fact: there exist renormalizable four-dimensional gauge theories that in the infrared regime generate the physics of gauge theories in $4+n$ dimensions below their cut-off scale M_s .¹ In that construction the Kaluza-Klein states of extra dimensions arise from having many gauge symmetries in four dimensions. The IR limit is determined by the dynamics of spontaneous breakdown of those symmetries.

The four-dimensional renormalizable theory constructed in ref. [1, 2] in the IR give the dynamics of gauge interactions in one extra dimension. Let us then first recall a few facts about non-abelian $SU(n)$ gauge theories in $4+1$ dimensions. They have three free parameters: the cut-off scale M_s , the dimensionful coupling constant

$$g_0 = 1/\sqrt{M} \equiv g_5/\sqrt{M_s} \quad (1)$$

and the compactification radius R (we assume that $M_s \gg 1/R$). The physics seen in four dimensions is obtained by integrating out the fifth dimension. The simplest compactifications are on a circle S_1 or on an orbifold S_1/Z_2 . In the latter case the four components of the vector potential $A_\mu(x_\mu, x_5)$, $\mu = 0, 1, 2, 3$, are even under Z_2 and the fifth component $A_5(x_\mu, x_5)$ is odd (here and in the following gauge group indices are suppressed). Fourier expanding in the discrete fifth component of momentum we define the four-dimensional

¹Identical construction has been independently given in ref. [2], except for explicit discussion of the interactions between the KK modes.

degrees of freedom (Kaluza-Klein modes): vectors $A_\mu^n(x_\mu)$, $n = 0, 1, \dots$, and scalars $A_5^n(x_\mu)$, $n = 1, \dots$. In the compactification on orbifold there is no zero mode of A_5 and, actually, $A_5(x_\mu, x_5)$ can be gauged away. If we compactify on a circle, the zero mode of A_5 remains in the physical spectrum (as a scalar in the adjoint representation of the $SU(n)$ group) and the massive vector KK modes are doubled. For the orbifold compactification, in the axial gauge $A_5 \equiv 0$, the effective Lagrangian after integrating over x_5 contains the following well known terms:

1. zero mode kinetic term invariant under $SU(n)$ gauge transformations in four dimensions

$$-\frac{1}{4}F_{\mu\nu}^{(0)}F^{(0)\mu\nu} \quad (2)$$

where

$$F_{\mu\nu}^{(0)} = \partial_\mu A_\nu^{(0)} - \partial_\nu A_\mu^{(0)} + \tilde{g}f A_\mu^{(0)} A_\nu^{(0)} \quad (3)$$

and $\tilde{g} = g_5/\sqrt{M_s R}$, with the coupling g_5 defined by eq.(1);

2. KK kinetic and mass terms

$$\sum_{n=1}^N (\partial_\mu A_\nu^n - \partial_\nu A_\mu^n)^2 + \sum_{n=1}^N \left(\frac{n\pi}{R}\right)^2 A_\mu^n A^{n\mu} \quad (4)$$

3. triple couplings

$$\begin{aligned} & \frac{2g}{\sqrt{M_s R}} f \sum_{n=1}^N \left[\partial_{[\mu} A_{\nu]}^0 A^{n\mu} A^{n\nu} + \partial_{[\mu} A_{\nu]}^n (A^{0\mu} A^{n\nu} + A^{n\mu} A^{0\nu}) \right] \\ & + \frac{g}{\sqrt{2M_s R}} f \sum_{n,m,l=1}^N \partial_{[\mu} A_{\nu]}^n A^{m\mu} A^{l\nu} \Delta_1(n, m, l) \end{aligned} \quad (5)$$

4. quartic couplings

$$\begin{aligned} & \frac{g^2}{M_s R} f f \sum_{n=1}^N \left(A_\mu^0 A_\nu^0 A^{n\mu} A^{n\nu} + \text{all permutations} \right) \\ & + \frac{g^2}{2M_s R} f f \sum_{n,m,l,k=1}^N A_\mu^n A_\nu^m A^{l\mu} A^{k\nu} \Delta_2(n, m, l, k) \end{aligned} \quad (6)$$

where the Δ_i are defined as:

$$\begin{aligned}\Delta_1 &= \delta(n+m-l) + \delta(n-m+l) + \delta(n-m-l) \\ \Delta_2 &= \delta(n+m-l-k) + \delta(n+m+l-k) + \delta(n+m-l+k) \\ &\quad + \delta(n-m+l+k) + \delta(n-m-l-k) + \delta(n-m+l-k) \\ &\quad + \delta(n-m-l+k).\end{aligned}\tag{7}$$

and f 's are the gauge group structure constants.

The structure of couplings reflects conservation of the fifth component of momentum. The truncation in the number of KK modes is understood, $n < N$ where $N \approx M_s R$, so that $M_N < M_s$. The effective Lagrangian has four-dimensional $SU(n)$ gauge invariance, with massive KK modes transforming linearly under the adjoint representation of $SU(n)$, but the full gauge invariance of the five-dimensional lagrangian is lost because of the truncation $n < N$.² The theory is manifestly non-renormalizable.

The renormalizable four-dimensional theory that, in its infrared region, generates the interactions described by the truncated Lagrangian (i-iv) is as follows [1, 2]. Let us consider the gauge structure

$$SU(n)_0 \times SU(n)_1 \times \dots SU(n)_N \equiv SU^{N+1}(n)\tag{8}$$

where the vector potentials are $A_{j\nu}^a$ and the dimensionless gauge coupling constants are equal for all of the $SU(n)$ symmetries, $g_{4j} \equiv g_4$. We suppose, in addition, that there is a set of scalar fields Φ_j ($j = 1, \dots, N$) (elementary or effective) which transform as $(\bar{n}, n)$ under $SU(n)_{j-1}$ and $SU(n)_j$ groups. We shall call Φ_j 's the link-Higgs fields. Their transformation properties define the links to be nearest-neighbour. The Lagrangian reads

$$\mathcal{L} = -\frac{1}{4} \sum_{i=0}^N F_{i\mu\nu}^a F^{a\mu\nu} + \sum_{i=1}^N (D_\mu \Phi)_i^\dagger D^\mu \Phi_i + V(\Phi_i)\tag{9}$$

²The Lagrangian for infinite tower of KK modes has full five-dimensional gauge invariance realized as non-linear gauge symmetries for the KK modes [6].

where $D_\mu = \partial_\mu + ig_4 \sum_{i=0}^N A_{i\mu}^a T_i^a$ (T_i^a are generators of the i th $SU(n)_i$ gauge symmetry), and the potential has full chiral symmetry $SU^N(n) \times SU^N(n)$ and is symmetric under interchange $\Phi_j \rightarrow \Phi_i$.

If the dynamics of the scalar sector (described here by the potential $V(\Phi_j)$; the scalars Φ_j can be replaced by technicolour-like condensates [2]) is such that the diagonal components of the scalars Φ_j acquire vacuum expectation value v , the chiral symmetry of the link-Higgs sector is spontaneously broken $SU^N(n) \times SU^N(n) \rightarrow SU^N(n)$ leaving N Nambu-Goldstone bosons, each transforming under the adjoint representation of $SU(n)$. The gauge symmetry $SU^{N+1}(n)$ of the full Lagrangian is also spontaneously broken to the diagonal $SU(n)$ and N vector bosons acquire masses eating up the Nambu-Goldstone bosons. If we want to have four-dimensional model corresponding to a compactification on a circle, we introduce one additional link-Higgs field Φ_0 transforming as $(3_0, \bar{3}_N)$. The chiral symmetry of the Higgs sector is now $SU^{N+1}(n) \times SU^{N+1}(n)$. Its breaking to $SU^{N+1}(n)$ leaves $N + 1$ Nambu-Goldstone bosons and N of them, as before, are eaten up by longitudinal components of N vector bosons. Thus, one massless scalar remains in the spectrum [3], in exact correspondence to the zero mode of A_5 in the 5d theory. The massless scalar transforms under the adjoint representation of the diagonal, unbroken, $SU(n)$ symmetry.

The physics of our renormalizable 4d theory below the scale v of spontaneous gauge symmetry breaking can be easily studied in detail in the non-linear σ -model approximation. For the orbifold compactification, upon substituting

$$\Phi_j \rightarrow v \exp(i\phi_j^a T^a / 2v) \quad (10)$$

the Φ_j kinetic terms lead to a mass matrix for the gauge fields

$$\sum_{i=1}^N \frac{1}{2} g_4^2 v^2 (A_{(i-1)\mu}^a - A_{i\mu}^a)^2 \quad (11)$$

This mass matrix has the structure of a nearest neighbour coupled oscillator

Hamiltonian. We can diagonalize the mass matrix to find the eigenvalues :

$$M_n = \sqrt{2}g_4v \sin \left[\frac{\gamma_n}{2} \right] \quad \gamma_n = \frac{n\pi}{N+1} , \quad n = 0, 1, \dots, N. \quad (12)$$

Thus we see that for small n this system has a KK tower of masses given by:

$$M_n \approx \frac{g_4v\pi n}{\sqrt{2}(N+1)} \quad n \ll N \quad (13)$$

and $n = 0$ corresponds to the zero-mode.

To match on to the spectrum of the KK modes, we require

$$\frac{g_4v}{\sqrt{2}(N+1)} = \frac{1}{R}. \quad (14)$$

Hence, the system with $SU(n)^{N+1}$ and $N \Phi_i$ provides a description of the KK modes of the 5d theory by generating the same mass spectrum.

Next, it is crucial to examine the interactions from the model. The gauge fields A_μ^j can be expressed as linear combinations of the mass eigenstates $\tilde{A}_\mu^n$ as:

$$A_\mu^j = \sum_{n=0}^N a_{jn} \tilde{A}_\mu^n. \quad (15)$$

The a_{nj} form a normalized eigenvector ($\vec{a}_n$) associated with the n th $n \neq 0$ eigenvalue and has the following components:

$$a_{nj} = \sqrt{\frac{2}{N+1}} \cos \left(\frac{2j+1}{2} \gamma_n \right) , \quad j = 0, 1, \dots, N, \quad (16)$$

The eigenvector for the zero-mode, $n = 0$, is always $\vec{a}_0 = \frac{1}{\sqrt{N+1}}(1, 1, \dots, 1)$. We can now rewrite the Lagrangian eqn.(9) in the mass eigenstates of the vector bosons ($\tilde{A}_\mu^n$) and derive the interactions between them. They are identical to those of the 5d theory, eq.(3-6), with the identification

$$\tilde{g} = \frac{g_5}{\sqrt{M_s R}} = \frac{g_4}{\sqrt{(N+1)}} \quad (17)$$

In the 4d theory, there are three relevant parameters, namely, the gauge coupling constant g_4 , the total number of $SU(3)$ groups $N+1$ and the VEV of

the Higgs field v determined by the potential $V(\Phi_i)$. The mappings between them and the parameters of the 5d theory are $N + 1 = M_s R$, $g_4 = g_5 = \sqrt{M_s/M}$ and $v = \sqrt{M_s M}$.

Another interesting fact is that the 4d model can also be interpreted as the 5d theory on a “transverse lattice” [4]. We construct a transverse lattice in the x^5 dimension where the lattice size is R and short-distance lattice cut-off is $a = 1/M_s$, so $N + 1 = R/a$. This is a foliation of $N + 1$ parallel branes, each spaced by a lattice cut-off a . On the i th brane we have an $SU(n)$ gauge theory denoted by $SU(n)_i$. The $SU(n)_i$ automatically have a common coupling constant g_4 .

The theory thus has N links in the x^5 direction that are continuous functions of x_μ . These correspond to the continuum limit Wilson lines:

$$\Phi_i(x^\mu) = \exp \left[i g_4 \int_{ia}^{(i+1)a} dx^5 A_5(x^\mu, x^5) \right] \rightarrow \exp \left[i g_4 a A_5 \left(x^\mu, (i + \frac{1}{2})a \right) \right] \quad (18)$$

The N Φ_i therefore transform as an $(\bar{n}, n)$ representation of $SU(n)_i \times SU(n)_{i+1}$ as in the 4d model. Φ_i is a unitary matrix and may be parameterized as in eq.(10).

The construction of ref. [1, 2] provides new insights and opens up several interesting possibilities. The most straightforward one is to discuss the 5d physics in the framework of a 4d renormalizable field theory. For instance, the power-law like running of the gauge couplings in 5 dimensions [5, 6] can be discussed in the renormalizable setting. In the 4d theory, the running is logarithmic but with the β function changing every time we pass the threshold M_n [1]. In the limit of large N one recovers the power-law like running.

Another obvious implication of these considerations is that the transverse lattice is *an exact* approach to 5d gauge theories with a cut-off.

Furthermore, one can use five-dimensional models as inspiration for four-dimensional theories and vice versa. Following the first direction, new ideas have been proposed on electroweak symmetry breaking [7, 8] and on supersymmetry breaking [9, 10] mechanisms in four dimensions. Going in the

reversed direction, one can get new insight into fermion mass generation in five dimensions [3]. A link to string theory is discussed in ref. [11].

Finally and perhaps most interestingly, the theory constructed in ref. [1, 2] offers an alternative, to the geometrical one, interpretation of gauge fields propagating in the bulk as arising from nearest-neighbour coupled, spontaneously broken, extended gauge symmetries in four dimensions. The big question is if similar interpretation can be extended to gravitational interactions. If so, one would get a new view on the notion of dimension.

Acknowledgments I thank P. Chankowski, J.P. Derendinger, E. Dudas, A. Falkowski, Ch. Grojean, Z. Lalak, H.P. Nilles and G. Ross for interesting conversations. This work was supported partially by the Polish State Committee for Scientific Research grant 5 P03B 119 20 for 2001-2002 and by the EC Contract HPRN-CT-2000-00152 for years 2000-2004.

References

- [1] C.T. Hill, S. Pokorski, J. Wang, hep-th/0104035 (to be published in *Phys. Rev. D*).
- [2] N. Arkani-Hamed, A.G. Cohen and H. Georgi, *Phys. Rev. Lett.* **86**, 4757 (2001), hep-th/0104005.
- [3] Hsin-Chia Cheng, C.T. Hill, S. Pokorski, J. Wang, preprint FERMILAB-PUB-01-053-T, hep-th/0104179 (to be published in *Phys. Rev. D*)
- [4] W.A. Bardeen, R.B. Pearson and E. Rabinovici, *Phys. Rev.* **D21**, 1037 (1980).
- [5] T.R. Taylor and G. Veneziano, *Phys. Lett.* **B212**, 147 (1988).
- [6] K.R. Dienes, E. Dudas and T. Gherghetta, *Nucl. Phys.* **B537**, 47 (1999);

- [7] N. Arkani-Hamed, A.G. Cohen and H. Georgi, preprint HUTP-01-A024, hep-ph/0105239.
- [8] H.-C. Cheng, Ch.T. Hill and J. Wang, preprint FERMILAB-PUB-01-079-T, hep-ph/0105323.
- [9] C. Csaki, J. Erlich, Ch. Grojean and G.D. Kribs, preprint UCB-PTH-01-18, hep-ph/0106044.
- [10] H.-C. Cheng, D.E. Kaplan, M. Schmalz and W. Skiba, preprint FERMILAB-PUB-01-108-T, hep-ph/0106098.
- [11] K. Sfetsos, preprint NEIP-01-005, hep-th/0106126.